

Week 6 workshop exercises

1. Find general solutions for the following differential equations:

$$(a) \frac{dy}{dx} = \frac{3x^2}{y} \qquad (b) \frac{dy}{dx} = 4xy^2$$

2. Solve the initial value problems for the following ODEs:

$$(a) \frac{dy}{dx} = \frac{y+1}{x-3}, \quad y(0) = 1 \qquad (b) \frac{dy}{dx} = \frac{x^2-1}{2y+1}, \quad y(0) = -1$$

3. Find the interval $\tau_{1/2}$ in which the amount of reactant A in a first order decay process

$$A(t) = A(0)e^{-kt}$$

is reduced by a factor of 2.

4. The following function was proposed as an approximation to $\cos(x)$ around $x = 0$:

$$f(x) = \frac{1 - \frac{115x^2}{252} + \frac{313x^4}{15120}}{1 + \frac{11x^2}{252} + \frac{13x^4}{15120}}$$

Using a calculator, inspect the performance of this approximation (a) at the origin; (b) in the interval between $-\pi/2$ and $+\pi/2$; (c) around $x = 2\pi$; (d) at infinity.

5. Using the substitution $x = kT/\Delta E$, demonstrate that the function

$$Q(T) = \frac{1}{2} + \frac{kT}{\Delta E} + \frac{\Delta E}{12kT}$$

can be an excellent approximation to the partition function of the ensemble of harmonic oscillators

$$Q(T) = \frac{1}{1 - e^{-\Delta E/kT}}$$

and use your calculator to estimate the interval of $kT/\Delta E$ for which the approximation remains valid.

6. Show that a differential equation of the form

$$\frac{dy}{dx} = f(ax + by + c)$$

where a, b, c are constants, is reduced to a separable form by the substitution $u = ax + by$. Proceed by calculating the differential du and using the result to eliminate dy from the numerator.