

CHEM1047 – Week 2 Lecture 2 – Well-behaved functions

□ Chapter 1, Sections 3-5 of Demidovich, "Problems in Mathematical Analysis", 2nd edition.

1. Discontinuities

At a simplified level, a *discontinuity* is a missing point or an abrupt jump in the function value from one point to the next. Discontinuities are common in mathematics, but rare in chemistry and physics for reasons that we will now discuss.




Figure 1. An example of a discontinuity in a function at the point where $x=0$.

Formally, a point of discontinuity of a function $f(x)$ is defined as a point x_0 where there is a difference between the right and the left limit of the function:

$$\lim_{x \rightarrow x_0^-} f(x) \neq \lim_{x \rightarrow x_0^+} f(x) \quad (1)$$

where the *left limit* $\lim_{x \rightarrow x_0^-} f(x)$ is defined as the value that the function approaches when x approaches x_0 from the left, and similarly for the *right limit*.

2. Restrictions imposed by physical reality

Let us calculate the energy required to create a discontinuity in the trajectory $x(t)$ of a particle of mass m . We will make a discontinuous function using the following limit:

$$x(t) = \lim_{k \rightarrow \infty} \left[\frac{1}{1 + e^{-kt}} \right] \quad (2)$$

It is clear from Figure 2 that the jump from $x=0$ to $x=1$ around $t=0$ gets sharper and sharper as the value of k increases. In the limit of infinite k the function has a discontinuity.




Figure 2. Time dependence the particle coordinate defined by Equation (2) when the parameter k is increased.

Let us now consider the velocity and the acceleration of the particle. To obtain the velocity, we must differentiate the coordinate with respect to time:

$$\frac{d}{dt} \left(\frac{1}{1+e^{-kt}} \right) = \frac{ke^{-kt}}{(1+e^{-kt})^2} \quad (3)$$

The result is plotted in Figure 3. As expected, larger values of k require greater velocities because the particle moves faster.




Figure 3. Time dependence the particle velocity defined by Equation (3) when the parameter k is increased.

After calculating the limit of the velocity for $k \rightarrow \infty$ at time zero

$$x'(t) = \lim_{k \rightarrow \infty} \left[\frac{ke^{-kt}}{(1+e^{-kt})^2} \right] \Rightarrow x'(0) = \lim_{k \rightarrow \infty} [k] = \infty$$

we conclude that *infinite velocity* would be required. Because energy is proportional to the square of the velocity, this means that *infinite energy* would have to be supplied to the particle to achieve a discontinuity in its trajectory. This is clearly impossible.

What about a sharp turn? The following function has a sharp corner when $k \rightarrow \infty$:

$$y(t) = \frac{1}{k} \ln [1 + e^{kt}] \quad (4)$$

Its plot for different values of k is shown in Figure 4.




Figure 4. Time dependence the particle coordinate defined by Equation (4) when the parameter k is increased.

Velocities are realistic here – the first derivative of $y(t)$ never goes to infinity:

$$y'(t) = \frac{e^{kt}}{1 + e^{kt}}$$

However, the *second derivative* now becomes infinite at $t = 0$ when $k \rightarrow \infty$:

$$y''(t) = \lim_{k \rightarrow \infty} \left[\frac{ke^{-kt}}{(1 + e^{-kt})^2} \right] \Rightarrow y''(0) = \lim_{k \rightarrow \infty} [k] = \infty$$

The second derivative of coordinate is acceleration. Newton's second law ($F = ma$) then means that *infinite force* would be required to create a sharp corner in the trajectory. That is also impossible.

Deeper inspection indicates that the same problem appears whenever there is a discontinuity in any of the derivatives of the state functions almost anywhere in chemistry and physics. It is therefore reasonable to create a category of functions that are admissible as realistic solutions to physical models. Such functions are called *well-behaved*. This is an informal term; it usually means: (a) finite; (b) continuous derivatives of any order; (c) finite integral; (d) finite integral of the square.

3. Formal definition of continuity

A function $f(x)$ defined in an interval (a, b) is called *continuous* in a point $x_0 \in (a, b)$ if

$$\lim_{x \rightarrow x_0} f(x) = f(x_0) \quad (5)$$

Sums, products and superpositions of continuous functions are continuous. Any function that is continuous in a particular interval is necessarily finite in that interval. If a function is continuous and *monotonic* in a certain interval, then its inverse function is also continuous and monotonic.

Examples of discontinuous functions:

1. Heaviside step function:

$$H(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases}$$

2. Signum function:

$$\text{sign}(x) = \begin{cases} +1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases}$$

3. Dirichlet function:

$$D(x) = \begin{cases} 0 & \text{if } x \text{ is an irrational number} \\ 1 & \text{if } x \text{ is a rational number} \end{cases}$$

Heaviside and signum functions are discontinuous in just one point; Dirichlet function is discontinuous everywhere because there are irrational numbers around every rational number and *vice versa*.

4. Discontinuities and singularities in physical theories

Note that the discussion above only applies to *solutions* of physical equations of motion. No such restrictions exist on the problem specifications – point particles, infinitely high walls and infinitely large masses are rather common as stage settings. A good example is Coulomb interaction energy between a pair of point charges, which goes to infinity as the inter-particle distance decreases:

$$E(r) = \frac{q_1 q_2}{4\pi\epsilon_0} \frac{1}{r}$$

The same applies to the gravitational potential of a point mass. A point at which a continuous function goes to infinity is called a *singularity*, and these are quite common in the *equations of motion*.




Figure 5. A plot of the absolute value of the gamma function of a complex argument.

A few singularities in the **gamma function** of a complex argument are shown in the figure above. In all cases, however, physically meaningful solutions to the equations of motion must be well-behaved. The only exception that appears to actually exist is **black holes**.